

## Swiss Run Greater Swiss Mountain Dogs When Too Spay or Neuter

When too spay or neuter is always the question. When is the right time? There really is no good answer for this question. Below is an article from Dr. Chris Zinc about spaying/neutering before or after a year of age. There are a number of pros and cons for spaying/neutering early, but the following discusses a few ideas on the subject.

### Orthopedic Considerations

“A study by Salmeri et al in 1991 found that bitches spayed at 7 weeks grew significantly taller than those spayed at 7 months, who were taller than those not spayed (or presumably spayed after the growth plates had closed).(1) A study of 1444 Golden Retrievers performed in 1998 and 1999 also found bitches and dogs spayed and neutered at less than a year of age were significantly taller than those spayed or neutered at more than a year of age.(2) The sex hormones, by communicating with a number of other growth-related hormones, promote the closure of the growth plates at puberty (3), so the bones of dogs or bitches neutered or spayed before puberty continue to grow. Dogs that have been spayed or neutered well before puberty can frequently be identified by their longer limbs, lighter bone structure, narrow chests and narrow skulls. This abnormal growth frequently results in significant alterations in body proportions and particularly the lengths (and therefore weights) of certain bones relative to others. For example, if the femur has achieved its genetically determined normal length at 8 months when a dog gets spayed or neutered, but the tibia, which normally stops growing at 12 to 14 months of age continues to grow, then an abnormal angle may develop at the stifle. In addition, with the extra growth, the lower leg below the stifle likely becomes heavier (because it is longer), and may cause increased stresses on the cranial cruciate ligament. In addition, sex hormones are critical for achieving peak bone density.(4) These structural and physiological alterations may be the reason why at least one recent study showed that spayed and neutered dogs had a higher incidence of CCL rupture.(5) Another recent study showed that dogs spayed or neutered before 5 1/2 months had a significantly higher incidence of hip dysplasia than those spayed or neutered after 5 1/2 months of age, although it should be noted that in this study there were no standard criteria for the diagnosis of hip dysplasia.(6) Nonetheless, breeders of purebred dogs should be cognizant of these studies and should consider whether or not pups they bred were spayed or neutered when considering breeding decisions.

### Cancer Considerations

A retrospective study of cardiac tumors in dogs showed that there was a 5 times greater risk of hemangiosarcoma, one of the three most common cancers in dogs, in spayed bitches than intact bitches and a 2.4 times greater risk of

hemangiosarcoma in neutered dogs as compared to intact males.(7) A study of 3218 dogs demonstrated that dogs that were neutered before a year of age had a significantly increased chance of developing bone cancer.(8) A separate study

showed that neutered dogs had a two-fold higher risk of developing bone cancer.(9) Despite the common belief that neutering dogs helps prevent prostate cancer, at least one study suggests that neutering provides no benefit.(10) There certainly is evidence of a slightly increased risk of mammary cancer in female dogs after one heat cycle, and for increased risk with each subsequent heat. While about 30 % of mammary cancers are malignant, as in humans, when caught and surgically removed early the prognosis is very good.(12) Luckily, canine athletes are handled frequently and generally receive prompt veterinary care.

### Behavioral Considerations

The study that identified a higher incidence of cranial cruciate ligament rupture in spayed or neutered dogs also identified an increased incidence of sexual behaviors in males and females that were neutered early.(5) Further, the study that identified a higher incidence of hip dysplasia in dogs neutered or spayed before 5 1/2 months also showed that early age neutering was associated with an increased incidence of noise phobias and undesirable sexual behaviors.(6) A recent report of the American Kennel Club Canine Health Foundation reported significantly more behavioral problems in spayed and neutered bitches and dogs. The most commonly observed behavioral problem in spayed females was fearful behavior and the most common problem in males was aggression.(12)

### Other Health Considerations

A number of studies have shown that there is an increase in the incidence of female urinary incontinence in dogs spayed early (13), although this finding has not been universal. Certainly there is evidence that ovarian hormones are critical for maintenance of genital tissue structure and contractility.(14, 15) Neutering also has been associated with an increased likelihood of urethral sphincter incontinence in males.(16) This problem is an inconvenience, and not usually life-threatening, but nonetheless one that requires the dog to be medicated for life. A health survey of several thousand Golden Retrievers showed that spayed or neutered dogs were more likely to develop hypothyroidism.(2) This study is consistent with the results of another study in which neutering and spaying was determined to be the most significant gender-associated risk factor for development of hypothyroidism.(17) Infectious diseases were more common in dogs that were spayed or neutered at 24 weeks or less as opposed to those undergoing neutering at more than 24 weeks.(18) Finally, the AKC-CHF report demonstrated a higher incidence of adverse reactions to vaccines in neutered dogs as compared to intact.(12)”

With everything said neutering every dog at or before the age of 6 months is not a black-and-white issue. Clearly more studies need to be done to evaluate the effects of spaying/neutering on young dogs. Granted, each dog is very individual, but overall waiting until a year of age is recommended.



Picture A 8 month old sisters



Picture B 8 month old sisters

Picture A: Girl on the left spayed at 6 months. Girl on the right not spayed yet.

Picture B: Girl on the left not spayed yet and Girl on the right spayed at 6 months.

#### References:

1. Salmeri KR, Bloomberg MS, Scruggs SL, Shille V.. Gonadectomy in immature dogs: effects on skeletal, physical, and behavioral development. *JAVMA* 1991;198:1193-1203
2. <http://www.grca.org/healthsurvey.pdf>
3. Grumbach MM. Estrogen, bone, growth and sex: a sea change in conventional wisdom. *J Pediatr Endocrinol Metab.* 2000;13 Suppl 6:1439-55.
4. Gilsanz V, Roe TF, Gibbens DT, Schulz EE, Carlson ME, Gonzalez O, Boechat MI. Effect of sex steroids on peak bone density of growing rabbits. *Am J Physiol.* 1988 Oct;255(4 Pt 1):E416-21.
5. Slauterbeck JR, Pankratz K, Xu KT, Bozeman SC, Hardy DM. Canine ovariohysterectomy and orchietomy increases the prevalence of ACL injury. *Clin Orthop Relat Res.* 2004 Dec;(429):301-5.
6. Spain CV, Scarlett JM, Houpt KA. Long-term risks and benefits of early-age gonadectomy in dogs. *JAVMA* 2004;224:380-387.
7. Ware WA, Hopper DL. Cardiac tumors in dogs: 1982-1995. *J Vet Intern Med* 1999 Mar-Apr;13(2):95-103
8. Cooley DM, Beranek BC, Schlittler DL, Glickman NW, Glickman LT, Waters D, *Cancer Epidemiol Biomarkers Prev.* 2002 Nov;11(11):1434-40
9. Ru G, Terracini B, Glickman LT. Host related risk factors for canine osteosarcoma. *Vet J.* 1998 Jul;156(1):31-9
10. Obradovich J, Walshaw R, Goullaud E. The influence of castration on the development of prostatic carcinoma in the dog. 43 cases (1978-1985). *J Vet Intern Med* 1987 Oct-Dec;1(4):183-7
11. [http://www.akcchf.org/pdfs/whitepapers/Biennial\\_National\\_Parent\\_Club\\_Canine\\_Health\\_Conference.pdf](http://www.akcchf.org/pdfs/whitepapers/Biennial_National_Parent_Club_Canine_Health_Conference.pdf)
12. Meuten DJ. *Tumors in Domestic Animals.* 4th Edn. Iowa State Press, Blackwell Publishing Company, Ames, Iowa, p. 575
13. Stocklin-Gautschi NM, Hassig M, Reichler IM, Hubler M, Arnold S. The relationship of urinary incontinence to early spaying in bitches. *J. Reprod. Fertil. Suppl.* 57:233-6, 2001
14. Pessina MA, Hoyt RF Jr, Goldstein I, Traish AM. Differential effects of estradiol, progesterone, and testosterone on vaginal structural integrity. *Endocrinology.* 2006 Jan;147(1):61-9.
15. Kim NN, Min K, Pessina MA, Munarriz R, Goldstein I, Traish AM. Effects of ovariectomy and steroid hormones on vaginal smooth muscle contractility. *Int J Impot Res.* 2004 Feb;16(1):43-50.
16. Aaron A, Eggleton K, Power C, Holt PE. Urethral sphincter mechanism incompetence in male dogs: a retrospective analysis of 54 cases. *Vet Rec.* 139:542-6, 1996
17. Panciera DL. Hypothyroidism in dogs: 66 cases (1987-1992). *J. Am. Vet. Med. Assoc.,* 204:761-7 1994
18. Howe LM, Slater MR, Boothe HW, Hobson HP, Holcom JL, Spann AC. Long-term outcome of gonadectomy performed at an early age or traditional age in dogs. *J Am Vet Med Assoc.* 2001 Jan 15;218(2):217-21.